

Design and implementation of bimolecular fluorescence complementation (BiFC) assays for the visualization of protein interactions in living cells

Tom K Kerppola

Howard Hughes Medical Institute and Department of Biological Chemistry, University of Michigan Medical School, Ann Arbor, Michigan 48109, USA. Correspondence should be addressed to T.K.K. (kerppola@umich.edu).

Published online 5 October; corrected online 30 November 2006 (details online); doi:10.1038/nprot.2006.201

Bimolecular fluorescence complementation (BiFC) analysis enables direct visualization of protein interactions in living cells. The BiFC assay is based on the discoveries that two non-fluorescent fragments of a fluorescent protein can form a fluorescent complex and that the association of the fragments can be facilitated when they are fused to two proteins that interact with each other. BiFC must be confirmed by parallel analysis of proteins in which the interaction interface has been mutated. It is not necessary for the interaction partners to juxtapose the fragments within a specific distance of each other because they can associate when they are tethered to a complex with flexible linkers. It is also not necessary for the interaction partners to form a complex with a long half-life or a high occupancy since the fragments can associate in a transient complex and un-associated fusion proteins do not interfere with detection of the complex. Many interactions can be visualized when the fusion proteins are expressed at levels comparable to their endogenous counterparts. The BiFC assay has been used for the visualization of interactions between many types of proteins in different subcellular locations and in different cell types and organisms. It is technically straightforward and can be performed using a regular fluorescence microscope and standard molecular biology and cell culture reagents.

INTRODUCTION

Studies of protein interactions have provided fundamental insights into the regulation of cellular functions. Many methods have been developed to investigate protein interactions. Most methods that enable direct detection of protein interactions, such as co-purification and affinity precipitation assays, require removal of the proteins from their native environment. By contrast, methods that enable detection of protein interactions in cells, such as genetic suppressor analysis, generally rely on indirect consequences of the protein interactions.

The visualization of protein interactions in living cells provides the potential for direct detection of protein interactions with minimal perturbation of their normal environment. Many strategies for the visualization of protein interactions in cells have been developed. The most commonly used are fluorescence resonance energy transfer (FRET)¹ and bimolecular fluorescence complementation (BiFC)². Others include fluorescence correlation spectroscopies^{3,4}, image correlation spectroscopy⁵ and complementation approaches using fragments of other proteins^{6–8}.

Comparison of BiFC and FRET

FRET and BiFC analysis are fundamentally different approaches and have complementary advantages and disadvantages. FRET is based on the transfer of excitation energy between two fluoro-

phores that are in close spatial proximity and have permissive relative orientations¹. This energy transfer results in a change in the fluorescence intensities and lifetimes of the two fluorophores. FRET analysis of a protein interaction requires quantification of the change in the fluorescence intensity or lifetime of the donor and the acceptor fluorophores in the presence versus the absence of energy transfer between the fluorophores.

BiFC is based on the association between fragments of a fluorescent protein when they are tethered in the same macromolecular complex (**Fig. 1**)². Thus, the association produces a fluorescent complex from non-fluorescent constituents. BiFC analysis requires determination of the difference in fluorescence intensities produced by the association of the fluorescent protein fragments in the presence versus the absence of an interaction between the proteins fused to the fragments. BiFC analysis has been used for the visualization of interactions between many different proteins in many cell types and organisms (see **Table 1**).

FRET enables, in principle, instantaneous monitoring of protein interactions whereas BiFC produces a signal after a delay required for the chemical reactions that generate the fluorophore. BiFC theoretically allows detection of interactions at lower protein concentrations and is predicted to be affected less by changes in cellular conditions that can alter the fluorescence intensities and lifetimes of fluorescent

Figure 1 | Schematic diagram representing the principle of the BiFC assay. Two fragments (YN and YC) of the yellow fluorescent protein (YFP) are fused to two putative interaction partners (A and B). An interaction between the proteins facilitates association between the fragments to produce a bimolecular fluorescent complex. The image on the right shows BiFC analysis of the recruitment of a protein to subnuclear foci by dimerization with a mutated version. Image on the right acquired by Nirmala Rajaram²⁵, reproduced with permission from the American Society for Microbiology.




TABLE 1 | Examples of protein interactions visualized using the BiFC assay

Category	Class	Proteins ^a	Organism	Reference
Peptides	Coiled coil	Anti-parallel NZ-CZ	<i>E. coli</i>	16
	Heat shock	Hsc70, Hsp90-TPR1, TPR2A, TPR2B	<i>E. coli</i>	17
Nuclear proteins	bZIP	Fos-Jun-ATF2; BATF-Jun; Maf-Sox	Mammalian cells	2, 23, 24, 68
	Rel	p50, IκB-p65	Mammalian cells	2
	bHLHZIP	Myc, Mxi1, Mad3, Mad4-Max, Mist-Mist	Mammalian cells	10, 25
	Bromodomain	AcH4-Brd2; SPA-1, P-TEFb-Brd4	Mammalian cells	26-28
	Smad	PKB/Akt, Smad4-Smad3	Mammalian cells	29
	IRF-Ets	IRF8-PU.1	Mammalian cells	31
	Winged helix	AcFKH1-CPCR1	<i>A. chrysogenum</i>	32
Ubiquitylation	E3 ligase-substrate	Skp2-Myc	Mammalian cells	22
		Grr1-Hof1	<i>S. cerevisiae</i>	33
		EID1-ASK1, 2, 4, 9, 13, 14, 15, SSK1 AtCUL1-ASK1, EID1, POCA-ASK1	<i>S. alba</i> , <i>P. crispum</i>	34, 35, 69
	Peptide conjugates	Jun-Ub, Jun-SUMO1	Mammalian cells	12
Plant pathogens	Type IV secretion	VirE2-VirD4	<i>A. tumefaciens</i>	36, 70
	Host-pathogen	VirE2, VirF, H2A-AtVIP, VirE3	<i>N. tabacum</i>	37, 38, 40
Signaling	MAP kinase network	MEKK3-IκBα; MEKK2-IκBβ; ERK1-p65; ERK2-p65	Mammalian cells	19
	PKB-PDK kinases	PKB/Akt, PDK1-hFt1	Mammalian cells	30
	Heterotrimeric G proteins	Gβ1-Gγ7	<i>D. discoideum</i> , Mammalian cells	41, 42
	Phospholipases	PLCβ2-PLCδ1	Mammalian cells	18
	Apoptosis	Bif1-Bax, TRAF6-Src	Mammalian cells	43, 64
	Photosensitivity	FpsA-FpsA	<i>A. nidulans</i>	44
Enzyme complexes	ACCS	ACS1, ACS4-ACS6, ACS7, ACS8	<i>E. coli</i>	45
	P450	P450C2, P450E1-P450 reductase; P450C2-BAP31	Mammalian cells	46, 47
Membrane proteins	Integrin signaling	Integrin αIibβ3, Syk-Src	Mammalian cells	48
	Arf GTPases	Arf1, Arf3, Arf4, Arf5-GBF1	Mammalian cells	49
	Lectin-glycoprotein	MCFD2, Cathepsin-ERGIC53	Mammalian cells	50
	Cytokine receptors	gp130-LIFR, gp130	Mammalian cells	51
	APP processing	APP-Notch2, APP	Mammalian cells	66
Nucleic acid binding	RNA binding	IMP, FMRP, hStau1, IRP1, PTB1-RNA; Nef-Nef; NXF1-Y14	Mammalian cells	52-54
	DNA binding	Zif268, PBSII-DNA	<i>In vitro</i>	71
Plant proteins	Transcription factors	FIE-MEA; bZIP63-bZIP63; LSD1-LSD1; bHLH1-OPF1; SAD, BPBF-GAMYB; LIP19-OBFI9; GRP23-RBP36B	<i>N. benthamiana</i> , <i>N. tabacum</i> , <i>A. thaliana</i> , <i>Allium sp.</i>	55-59, 63
		PFTα-PFTβ; T143c-T143c	<i>N. benthamiana</i> , <i>A. thaliana</i>	55, 56
	Flowering	FD-FT	<i>N. benthamiana</i>	60
	Plastid division	MinD1-MinE1; FtsZ1, ARC6-FtsZ2	<i>N. tabacum</i>	61
	Enzyme complex	AtSufE-AtSufS, AtNifS	<i>N. tabacum</i>	65

^aProtein pairs that have been tested are separated by a dash. In cases where several protein pairs have been tested, the alternative partners are separated by a comma. Different combinations of proteins that have been tested are separated by semi-colons.

proteins. FRET requires close spatial proximity between the fluorophores whereas BiFC requires that the fluorescent protein fragments have the dynamic flexibility to associate in the protein complex. FRET requires that a large fraction of the fluorescent proteins

associate with each other whereas BiFC can detect an interaction involving only a small subset of each fusion protein. Each approach is therefore applicable for different purposes. The fundamental principles and recent applications of the BiFC assay have been reviewed⁹.

Applications of the BiFC assay

The BiFC assay is applicable for visualization of the steady-state distributions of complexes formed between virtually any combination of proteins in a wide variety of cell types and organisms. Although the approach is applicable for studies of protein interactions in a wide variety of organisms (see **Table 1**), the chemistry of fluorophore formation requires molecular oxygen, making this approach unsuitable for use in organisms that are obligate anaerobes. The proteins must be able to accept fusions to fluorescent protein fragments without disruption of their functions. The complex must tolerate stabilization of the interaction by the association between the fluorescent protein fragments without changes in function or deleterious effects for the cell. The BiFC assay is particularly valuable for determining the subcellular locations of protein interactions, which can provide insight into the biological functions of newly discovered protein complexes. Fragments of several proteins that can associate when they are brought together by an interaction between proteins fused to the fragments have been identified. These include fragments of ubiquitin, β -galactosidase and dihydrofolate reductase^{6–8}. The advantage of BiFC analysis is that association between fragments of fluorescent proteins produces a complex with intrinsic fluorescence, eliminating the need for exogenous stains and enabling direct detection of the protein complex. We have developed several extensions of the BiFC assay that enable visualization of multiple protein interactions in the same cell^{10,11} as well as covalent protein modifications¹². Recent studies have also suggested that BiFC analysis can be used to determine the topology of membrane proteins¹³ and for high-throughput screening for the effects of small molecules on protein complexes¹⁴. Future efforts will undoubtedly identify many new applications for fluorescence complementation.

Limitations of the BiFC assay

The BiFC assay has several characteristics that limit its applicability and should temper interpretations based on results from this assay. One limitation of the BiFC approach is the time required for fluorophore maturation, which reflects the chemical reactions required for formation of the cyclic fluorophore. This prevents real-time detection of rapid changes in interactions using the BiFC assay. It is possible that some of the chemical reactions required for fluorophore formation occur in the isolated fragments, accelerating fluorescence complementation under some conditions¹⁵.

Bimolecular fluorescent complex formation is also likely to affect the dynamics of complex dissociation and partner exchange. Under many *in vitro* conditions, formation of the bimolecular fluorescent

complex is essentially irreversible^{2,16,17}. However, in some experiments, rapid changes in the fluorescence signal have been observed^{18,19}. It is possible that bimolecular fluorescent complex formation is reversible under these conditions, but it is difficult to exclude the possibility that other processes, such as protein degradation, affect the signal in living cells. The signal from fluorescence complementation has also been shown to decrease under conditions predicted to reduce complex formation under some *in vitro* conditions^{15,18}. However, dissociation of the fusion proteins has not been directly demonstrated.

Finally, fluorescent protein fragments have a finite ability to associate with each other independent of an interaction between proteins fused to the fragments. This major source of background signal in the BiFC assay varies depending on the identities of the fusion proteins and their levels of expression. Generally, this problem can be alleviated by expression of the fusion proteins at concentrations approximating their endogenous counterparts. In cases where this may not be possible, alternative fusions can be tested for the specificity of fluorescence complementation.

Strengths of the BiFC assay

There are many characteristics of the BiFC assay that make it useful for the study of protein interactions. First, it enables direct visualization of protein interactions and does not rely on their secondary effects. Second, the interactions can be visualized in living cells, eliminating potential artifacts caused by cell lysis or fixation. Third, the proteins are expressed in a relevant biological context, ideally at levels comparable to those of their endogenous counterparts. This increases the likelihood that the results reflect the properties of native proteins, including potential effects of post-translational modifications. Fourth, the BiFC assay does not require stoichiometric complex formation but can be used to detect interactions between subpopulations of each protein. Finally, BiFC does not require specialized equipment, apart from an inverted fluorescence microscope equipped with objectives that allow imaging of fluorescence in cells. The direct detection of bimolecular complex formation requires no post-acquisition image processing for interpretation of the data.

This protocol focuses on the visualization of protein interactions in cultured mammalian cells using the BiFC assay², but the general principles described are applicable to many other experimental systems. Although this protocol describes the use of transiently transfected plasmid vectors, it is likely that other expression strategies (retroviruses, lentiviruses, etc.) can be used with only slight modifications to the protocol.

MATERIALS

REAGENTS

- Plasmid vectors for expression of fusion proteins
- DNA encoding N-terminal fragment of fluorescent protein (see **Table 2**)
- DNA encoding C-terminal fragment of fluorescent protein (see **Table 2**)
- DNA encoding proteins of interest (interaction partners, wild type)
- DNA encoding mutated proteins of interest that do not interact with each other (negative controls)
- Cells that can be transfected with plasmid DNA (preferably adherent, monolayer cell line)
- Materials for cell culture
- Materials for cell transfection

- Tissue culture vessels appropriate for experiment: e.g., cluster plates, slide chambers, or glass coverslips
- Additional reagents and equipment for expressing proteins in mammalian cells, mammalian cell culture, and immunoblotting

EQUIPMENT

- Inverted fluorescence microscope
- Sensitive charge-coupled device (CCD) camera
- 20 \times to 100 \times microscope objectives
- Microscope filters for visualization of fluorescent proteins: YFP, Venus (excitation, 500 \pm 10 nm; emission, 535 \pm 15 nm) and CFP, Cerulean (excitation, 436 \pm 5 nm; emission, 470 \pm 15 nm)
- Software for instrument control and data analysis
- Facilities for cell culture

TABLE 2 | Combinations of fluorescent protein fragments recommended for BiFC analysis

Fusions ^a	Purpose	Excitation filter	Emission filter	Reference
A–YN155	A–B interaction	500/20 nm	535/30 nm	2
B–YC155				
A–YN173	A–B interaction	500/20 nm	535/30 nm	11
B–YC173				
A–VN155	A–B interaction	500/20 nm	535/30 nm	21
B–VC155				
A–VN173	A–B interaction	500/20 nm	535/30 nm	21
B–VC173				
A–CN155	A–B interaction	436/10 nm	470/30 nm	11
B–CC155				

^aA and B correspond to proteins whose interaction is to be tested. YN155 corresponds to residues 1–154 of EYFP⁷². YC155 corresponds to residues 155–238 of EYFP. YN173 corresponds to residues 1–172 of EYFP. YC173 corresponds to residues 173–238 of EYFP. VN155 corresponds to residues 1–154 of Venus²⁰. VC155 corresponds to residues 155–238 of Venus. VN173 corresponds to residues 1–172 of Venus. VC173 corresponds to residues 173–238 of Venus. CN155 corresponds to residues 1–154 of ECFP⁷². CC155 corresponds to residues 155–238 of ECFP.

PROCEDURE


1 | Select the fluorescent protein fragments to be used. Several combinations of fluorescent protein fragments support bimolecular fluorescence complementation¹¹; those recommended for BiFC analysis are listed in **Table 2**. For most purposes, fragments of YFP truncated at residue 155 (designated YN155 and YC155) are recommended, because they exhibit a relatively high complementation efficiency when fused to many interaction partners, yet produce low fluorescence when fused to proteins that do not interact with each other². Fragments of YFP truncated at residue 173 (designated YN173 and YC173) can also be used¹¹, and may exhibit a different efficiency of complementation owing to differences in the steric constraints imposed by tethering of the fragments to the protein complex. Fragments of Venus (a mutated GFP with high fluorescence intensity)²⁰ truncated at either residue 155 or 173 (designated VN155 and VC155, or VN173 and VC173, respectively) produce a significantly brighter fluorescent signal when fused to specific interaction partners²¹. However, these fragments also produce a brighter signal when fused to proteins that do not selectively interact with each other²¹. These fragments have the great advantage that the bimolecular fluorescent complex is readily detectable at 37 °C, which avoids the incubation at 30 °C that is generally necessary to detect complementation using YFP fragments. Other combinations of fluorescent protein fragments can also be used, especially when using BiFC analysis for the visualization of multiple protein complexes in the same cell¹¹.

2 | Determine the sites where the fluorescent protein fragments can be fused to the putative interaction partners. Determine the positions of the fusions empirically to fulfill the three criteria described below.

First, ensure that the fusions allow the fragments of the fluorescent proteins to associate with each other if the putative partners interact. Information about the structure and location of the interaction interface may be useful to determine optimal positions for the fusions. However, this information is not essential because fusions that can be used for BiFC analysis can be identified by screening multiple combinations of fusion proteins for fluorescence complementation. One strategy for the identification of fusion proteins that allow bimolecular fluorescence complementation is to fuse each of the fluorescent protein fragments to the N- and C-terminal end of each interaction partner, and to test for complementation in all eight combinations that contain both fragments of the fluorescent protein (**Fig. 2**).

Second, confirm that fusions do not affect the localization or the stabilities of the proteins by comparing the localization and expression levels of the fusion proteins with those of wild-type proteins lacking the fusions; indirect immunofluorescence and immunoblot analyses can be used.

Figure 2 | Combinations of fusion proteins to be tested for bimolecular fluorescence complementation. Two fragments (YN and YC) of the yellow fluorescent protein (YFP) are fused to two putative interaction partners (A and B). Fusion proteins that produce optimal signal must generally be empirically determined. Multiple combinations of fusion proteins should be tested for bimolecular fluorescence complementation. Amino- and carboxyl-terminal fusions can be used to test eight distinct combinations (**a–h**). Although it may appear that combinations **e–h** might not be favorable for bimolecular complex formation, this will depend on the precise structures and flexibilities of the fusion proteins, which are difficult to predict. For true interaction partners, it is virtually always possible to find fusion proteins that produce a detectable signal.



Third, test the fusion proteins for all known functions of the endogenous proteins to ensure that the fusions do not affect the functions of the proteins under investigation.

? TROUBLESHOOTING

3| Select linkers to connect the fragments to the proteins of interest. The linkers must provide flexibility for independent motion of the fluorescent protein fragments and the interaction partners, allowing the fragments to associate when the proteins interact. We have used the RSIAT and RPACKIPNDLKQKVMNH linker sequences in many fusion constructs used for BiFC analysis^{2,11}. These linkers have been used for the visualization of interactions between many structurally unrelated proteins. The sequence AAANSSIDLISVPVDSR encoded by the multiple cloning sites of the pCMV-FLAG vector (Sigma) has also been successfully used as a linker in many BiFC experiments. A peptide sequence designed to be flexible, such as (GGGS)_n, can also be used, although it can potentially affect the degradation of the fusion protein. Although these linker sequences have worked well for the proteins examined previously, it is possible that linkers of a different length or sequence are optimal for BiFC analysis of interactions between other proteins.

? TROUBLESHOOTING

4| Select a cell culture system. Choose a cell culture system that represents the biological context to be investigated, and allows efficient introduction of DNA into a large fraction of the cells. Cells that grow as an adherent monolayer are generally easier to image. The BiFC assay has been used for the analysis of protein interactions in many mammalian cell lines including COS-1, HEK293, HeLa, Hep3B, α TN4, and NIH3T3 cells as well as in intact organisms^{2,10,12,18,19,22–66}.

5| Select a strategy for expression of the fusion proteins. Choose either transient expression (A) or stable expression (B) strategies, based on the purpose of the experiment.

(A) Transient overexpression of the fusion proteins


This approach may be adequate to determine if a pair of proteins can interact in cells and to determine the subcellular location of the complex. To minimize protein mislocalization and formation of non-native complexes as a result of overexpression, express the fusion proteins at levels comparable to the endogenous proteins. This can be achieved by using plasmids with weak promoters, by transfecting small amounts of plasmid DNA and by observing the cells as soon as signal is detectable.

(B) Expression in stable cell lines

More reproducible levels of expression can be obtained by using inducible expression vectors integrated into the genomes of stable cell lines. This allows for the control of protein expression at relatively uniform levels in the entire cell population and replication of experiments at constant expression levels, independent of transfection efficiency and other factors that are difficult to control in transient assays.

6| Design controls to determine whether complementation reflects a specific protein interaction. As fluorescent protein fragments are able to form fluorescent complexes with a low efficiency in the absence of a specific interaction, it is essential to include negative controls in each experiment. Spontaneous complementation is generally reduced when the fragments are fused to proteins that do not interact with each other; appropriate negative controls are fusion proteins in which the interaction interface has been mutated and fused to the fluorescent protein fragments in a manner identical to the wild-type fusion proteins (Fig. 3)^{2,10}. Compare the level of expression and localization of the mutated and wild-type fusion proteins by immunoblot and indirect immunofluorescence analyses (using any standard or commercially available method). Quantify and compare the efficiencies of fluorescence complementation between the wild-type and mutated proteins. If there is no prior knowledge of the location or the structural nature of the interaction interface, it is possible to screen for mutations that alter the efficiency of bimolecular fluorescence complementation, and thereby determine if the complementation reflects a specific interaction.

Figure 3 | Determination of the specificity of bimolecular fluorescence complementation by mutational analysis. The requirement for a specific protein interaction for fluorescence complementation should be tested by examining the effects of mutations that prevent the association of the interaction partners (data adapted from ref. 2, reproduced with permission from Elsevier). The wild type (a) and mutated (b) interaction partners should be expressed at equal concentrations together with an internal reference (i.e., CFP). The fluorescence intensities produced by bimolecular fluorescence complementation (YN–YC) and the internal reference (CFP) are measured in individual cells. The distribution of ratios between the fluorescence intensities in individual cells is plotted in each histogram. The colored bars indicate cells with different ratios of YN–YC to CFP fluorescence. When the proteins interact (a), the fluorescence intensity of YN–YC will increase, resulting in a higher fluorescence ratio. The interaction may occur in all or in a subset of cells. When no interaction is observed (b), most cells will have a very low fluorescence ratio.



The BiFC assay can therefore be used to determine whether two proteins interact in cells without prior knowledge of the location or the structural nature of the interaction interface.

Use some of the numerous fusion proteins whose interactions have been visualized using the BiFC assay as positive controls (see **Table 1**). However, the failure to detect fluorescence complementation between the proteins under investigation does not demonstrate the absence of an interaction (see ANTICIPATED RESULTS).

7| Prepare plasmids encoding fusion proteins. Construct plasmid expression vectors, using the appropriate vectors, by fusing the sequences encoding the selected fluorescent protein fragments (see **Table 2**) to the sequences encoding the proteins of interest. Any standard cloning techniques can be used. Whenever possible, test fusions to both the N- and C-terminal ends of the proteins to be investigated (**Fig. 2**). Construct negative control plasmids that encode mutated non-interacting variants of the proteins (see Step 6) using the same strategy. Positive controls should be included to ensure that a known interaction can be detected (see Step 6).

? TROUBLESHOOTING

8| Prepare cells for transfection. Seed cells the day before transfection at an appropriate density. This density should allow for cell proliferation over the course of the experiment while taking into consideration the effects of cell growth and density on the interaction under investigation. Cluster plates are convenient for processing multiple transfections in parallel. If short-working-distance objectives will be used to visualize the interaction, grow the cells in slide chambers or on glass coverslips.

? TROUBLESHOOTING

9| Transfect cells. Transfect cells (using the optimal procedure for the cells) with appropriate amounts (e.g., 0.25 µg) of the BiFC plasmids encoding the fusion proteins when an appropriate confluency (~50%) is reached. In parallel, transfect cells with the negative and positive control plasmids. For quantification of the efficiency of fluorescence complementation, all plasmids should be co-transfected with the same amount of an internal control plasmid (e.g., a plasmid that expresses CFP, see **Fig. 3**).

? TROUBLESHOOTING

10| Allow time for fusion protein expression and fluorophore maturation. Grow cells under conditions appropriate for the cell type until fluorescence is detected (12–36 h). If necessary, incubate the cells at 30 °C with 5% CO₂ to promote maturation of the fluorophore and to increase the signal.

! CAUTION Results obtained under low temperature conditions should be interpreted with care, because incubation at a lower temperature could alter protein interactions.

? TROUBLESHOOTING

11| Remove dead cells. Wash the cells once with an amount of PBS sufficient to remove dead cells and cell debris, then add fresh medium.

12| Observe cells. Image the cells using an inverted fluorescence microscope. When using cells grown on plastic, a long-working-distance objective is convenient, but produces lower signal (owing to the lower numerical aperture) than a short-working-distance objective. A 20× objective is useful for observing large numbers of cells and can provide general subcellular localization information, whereas 60× or 100× objectives can be necessary for detailed localization within subcellular compartments. For detection of complementation between YFP or Venus fragments an excitation filter with 500 ± 10 nm transmission and an emission filter with 535 ± 15 nm transmission are appropriate. Confirm that fluorescent cells are alive by comparing their morphology with that of non-transfected cells. Cells grown on coverslips can be fixed and individual proteins can be visualized by indirect immunofluorescence analysis. Use protocols for fixation and immunofluorescence that have been established for the cell line and antibody to be used.

? TROUBLESHOOTING

13| Establish the levels of protein expression. Compare the levels of fusion protein expression with those of the endogenous proteins by immunoblot analysis. The cultures used for imaging can be subsequently processed for immunoblotting or separate cultures can be prepared in parallel. Use protocols for immunoblotting that have been established to work for the cell line and antibody to be used. Use the ratio between the intensities of the bands corresponding to the transfected and endogenous proteins together with the transfection efficiency to estimate the relative levels of transfected and endogenous proteins. Ideally, the amount of transfected protein should not exceed that of the endogenous protein in the cells.

? TROUBLESHOOTING

14| Analyze the data. Compare the intensities and numbers of fluorescent cells observed when the cells are transfected with the wild-type interaction partners with those observed when the cells are transfected with the negative control constructs. For quantitative analysis of the efficiencies of fluorescence complementation, divide the fluorescence intensities produced by



PROTOCOL

fluorescence complementation by the fluorescence intensities produced by intact fluorescent protein in individual cells (see **Fig. 3**). Higher fluorescence intensity and an increased number of fluorescent cells for the wild-type proteins is consistent with a specific interaction.

● TIMING

The construction of plasmid vectors for the expression of fusion proteins can be accomplished in a few days once the design for the vectors has been completed. Preparation of cells that express the fusion proteins can vary a great deal from days in transient transfection experiments to months or years for the production of stable cell lines or transgenic organisms. In transient expression experiments, fluorescence from specific interactions can generally be detected between 12 and 30 h after transfection. In the case of complementation between YFP and CFP fragments, this generally requires a short incubation at 30 °C to facilitate fluorophore maturation. In the case of complementation between Venus fragments, this step is generally not necessary. Longer incubation should be avoided because this may result in higher expression of fusion proteins and complementation due to nonspecific interactions. Images can be recorded and analyzed in less than 1 h for each combination of fusion proteins.

? TROUBLESHOOTING

See **Table 3** for troubleshooting advice.

TABLE 3 | Troubleshooting table.

STEPS	PROBLEM	SOLUTION
7, 8, 9	Cell culture, transfection	Problems with cell culture and transfection will prevent detection of protein interactions. Use plasmids encoding intact fluorescent proteins to establish the transfection efficiency and expression of heterologous proteins in the cells.
10, 13	Protein expression	Problems with protein expression can be generic or specific to the fusion proteins to be tested. Establish that the chosen expression vectors are appropriate for the cells that are used. Determine the expression of the fusion proteins by western blot analysis. If the fusion proteins cannot be detected, construct new expression vectors with a different arrangement of fluorescent protein fragments or change the linker sequences.
2, 13	Protein localization	If the fusion proteins have a different subcellular distribution than the endogenous proteins, it is possible that overexpression of the proteins causes their mislocalization. Reduce the level of fusion protein expression by using regulated expression vectors, ideally as stably transfected integrants. If the proteins are transiently expressed, shorten the time allowed for protein expression or reduce the amount of plasmid transfected into the cells. It is also possible that the fusions or the interaction between them affect the distributions of the proteins. Test different arrangements of the fluorescent protein fragments and compare the distributions of the fusion proteins when expressed singly or in combination.
2, 3, 12	Complementation	If the proteins are expressed but no fluorescence complementation is detected, it is possible that the steric arrangement of fluorescent protein fragments does not allow their association. Test other arrangements of fluorescent protein fragments or change the linker sequences. If the cells are grown at 37 °C, transfer them to 30 °C for a few hours before imaging to promote fluorophore maturation. It is also possible that the proteins do not interact under the conditions tested.

ANTICIPATED RESULTS

Interpretation of results from BiFC analysis

If fluorescence is detected when wild-type proteins are expressed, and this signal is eliminated or significantly reduced by single amino acid substitutions that prevent the interaction, it is likely that the bimolecular fluorescence complementation represents a specific interaction between the proteins fused to the fragments of fluorescent proteins.

If mutations that are known to eliminate the interaction do not reduce or eliminate the fluorescence, then the bimolecular complementation is due to a nonspecific interaction between the fusion proteins. If this is the case, the BiFC assay may not be an appropriate assay for the study of the two proteins. If the levels of expression of the fusion proteins were higher than those of their endogenous counterparts, modify the protocol or use another expression system to achieve levels of protein expression more consistent with those of the endogenous proteins. Alternatively, a different combination of fusion proteins or linkers should be tested.

If no fluorescence complementation is detected in the BiFC assay it does not prove the absence of an interaction, even if coexpression of the same fusion proteins with other interaction partners results in bimolecular fluorescence complementation. Different combinations of BiFC constructs or alternative approaches should be employed. Fluorescence in a small subpopulation of cells is difficult to interpret, since it may represent complementation due to nonspecific interactions, or it may reflect a regulated interaction that occurs only in response to signaling in a subset of cells. Further studies are necessary to distinguish these possibilities.

The fluorescence intensity produced by bimolecular fluorescence complementation in living cells is generally less than 10% of that produced by intact fluorescent proteins. It is likely that only a subset of the fragments associate with each other because the fluorescence intensity of BiFC complexes produced *in vitro* is comparable with that of intact fluorescent proteins². Several variants of the BiFC assay have been developed that enable visualization of multiple protein interactions in the same cell as well as covalent protein modifications^{9,11,67}.

ACKNOWLEDGMENTS I thank C.-D. Hu for his participation in the design and implementation of the BiFC assay in mammalian cells and all members of the Kerppola laboratory for their contributions to the improvement and adaptation of the BiFC approach.

COMPETING INTERESTS STATEMENT The author declares that he has no competing financial interests.

Published online at <http://www.natureprotocols.com>

Rights and permissions information is available online at <http://npg.nature.com/reprintsandpermissions>

- Förster, T. 10th Spiers Memorial Lecture. Transfer mechanisms of electronic excitation. *Discuss. Faraday Soc.* **27**, 7–17 (1959).
- Hu, C.D., Chinenov, Y. & Kerppola, T.K. Visualization of interactions among bZIP and Rel family proteins in living cells using bimolecular fluorescence complementation. *Mol. Cell* **9**, 789–798 (2002).
- Brock, R. & Jovin, T.M. Fluorescence correlation microscopy (FCM)-fluorescence correlation spectroscopy (FCS) taken into the cell. *Cell. Mol. Biol.* **44**, 847–856 (1998).
- Baudendistel, N., Muller, G., Waldeck, W., Angel, P. & Langowski, J. Two-hybrid fluorescence cross-correlation spectroscopy detects protein-protein interactions *in vivo*. *Chemphyschem* **6**, 984–990 (2005).
- Petersen, N.O., Hodelius, P.L., Wiseman, P.W., Seger, O. & Magnusson, K.E. Quantitation of membrane receptor distributions by image correlation spectroscopy: concept and application. *Biophys. J.* **65**, 1135–1146 (1993).
- Johnsson, N. & Varshavsky, A. Split ubiquitin as a sensor of protein interactions *in vivo*. *Proc. Natl. Acad. Sci. USA* **91**, 10340–10344 (1994).
- Rossi, F., Charlton, C.A. & Blau, H.M. Monitoring protein-protein interactions in intact eukaryotic cells by beta-galactosidase complementation. *Proc. Natl. Acad. Sci. USA* **94**, 8405–8410 (1997).
- Pelletier, J.N., Campbell-Valois, F.X. & Michnick, S.W. Oligomerization domain-directed reassembly of active dihydrofolate reductase from rationally designed fragments. *Proc. Natl. Acad. Sci. USA* **95**, 12141–12146 (1998).
- Kerppola, T.K. Visualization of molecular interactions by fluorescence complementation. *Nat. Rev. Mol. Cell Biol.* **7**, 449–456 (2006).
- Grinberg, A.V., Hu, C.D. & Kerppola, T.K. Visualization of Myc/Max/Mad family dimers and the competition for dimerization in living cells. *Mol. Cell. Biol.* **24**, 4294–4308 (2004).
- Hu, C.D. & Kerppola, T.K. Simultaneous visualization of multiple protein interactions in living cells using multicolor fluorescence complementation analysis. *Nat. Biotechnol.* **21**, 539–545 (2003).
- Fang, D.Y. & Kerppola, T.K. Ubiquitin-mediated fluorescence complementation reveals that Jun ubiquitinated by Itch/AIP4 is localized to lysosomes. *Proc. Natl. Acad. Sci. USA* **101**, 14782–14787 (2004).
- Zamyatnin, A.A. *et al.* Assessment of the integral membrane protein topology in living cells. *Plant J.* **46**, 145–154 (2006).
- MacDonald, M.L. *et al.* Identifying off-target effects and hidden phenotypes of drugs in human cells. *Nat. Chem. Biol.* **2**, 329–337 (2006).
- Demidov, V.V. *et al.* Fast complementation of split fluorescent protein triggered by DNA hybridization. *Proc. Natl. Acad. Sci. USA* **103**, 2052–2056 (2006).
- Ghosh, I., Hamilton, A.D. & Regan, L. Antiparallel leucine zipper-directed protein reassembly: application to the green fluorescent protein. *J. Am. Chem. Soc.* **122**, 5658–5659 (2000).
- Magliery, T.J. *et al.* Detecting protein-protein interactions with a green fluorescent protein fragment reassembly trap: Scope and mechanism. *J. Am. Chem. Soc.* **127**, 146–157 (2005).
- Guo, Y.J., Rebecchi, M. & Scarlata, S. Phospholipase C beta(2) binds to and inhibits phospholipase C delta(1). *J. Biol. Chem.* **280**, 1438–1447 (2005).
- Schmidt, C. *et al.* Mechanisms of proinflammatory cytokine-induced biphasic NF-kappa B activation. *Mol. Cell* **12**, 1287–1300 (2003).
- Nagai, T. *et al.* A variant of yellow fluorescent protein with fast and efficient maturation for cell-biological applications. *Nat. Biotechnol.* **20**, 87–90 (2002).
- Shyu, Y.J., Liu, H., Deng, X. & Hu, C.D. Identification of new fluorescent protein fragments for bimolecular fluorescence complementation analysis under physiological conditions. *Biotechniques* **40**, 61–66 (2006).
- von der Lehr, N. *et al.* The F-Box protein Skp2 participates in c-Myc proteasomal degradation and acts as a cofactor for c-Myc-regulated transcription. *Mol. Cell* **11**, 1189–1200 (2003).
- Deppmann, C.D., Thornton, T.M., Utama, F.E. & Taparowsky, E.J. Phosphorylation of BATF regulates DNA binding: a novel mechanism for AP-1 (activator protein-1) regulation. *Biochem. J.* **374**, 423–431 (2003).
- Rajaram, N. & Kerppola, T.K. Synergistic transcription activation by Maf and Sox and their subnuclear localization are disrupted by a mutation in Maf that causes cataract. *Mol. Cell. Biol.* **24**, 5694–5709 (2004).
- Zhu, L.Q. *et al.* Inhibition of Mist1 homodimer formation induces pancreatic acinar-to-ductal metaplasia. *Mol. Cell. Biol.* **24**, 2673–2681 (2004).
- Kanno, T. *et al.* Selective recognition of acetylated histones by bromodomain proteins visualized in living cells. *Mol. Cell* **13**, 33–43 (2004).
- Farina, A. *et al.* Bromodomain protein Brd4 binds to GTPase-activating SPA-1, modulating its activity and subcellular localization. *Mol. Cell. Biol.* **24**, 9059–9069 (2004).
- Jang, M.K. *et al.* The bromodomain protein Brd4 is a positive regulatory component of P-TEFb and stimulates RNA polymerase II-dependent transcription. *Mol. Cell* **19**, 523–534 (2005).
- Remy, I., Montmarquette, A. & Michnick, S.W. PKB/Akt modulates TGF-beta signalling through a direct interaction with Smad3. *Nat. Cell Biol.* **6**, 358–365 (2004).
- Remy, I. & Michnick, S.W. Regulation of apoptosis by the Ft1 protein, a new modulator of protein kinase B/Akt. *Mol. Cell. Biol.* **24**, 1493–1504 (2004).
- Laricchia-Robbio, L. *et al.* Partner-regulated interaction of IFN regulatory factor 8 with chromatin visualized in live macrophages. *PNAS* **102**, 14368–14373 (2005).
- Hoff, B. & Kuck, U. Use of bimolecular fluorescence complementation to demonstrate transcription factor interaction in nuclei of living cells from the filamentous fungus *Acremonium chrysogenum*. *Curr. Genet.* **47**, 132–138 (2005).
- Blondel, M. *et al.* Degradation of Hof1 by SCF^{Grr1} is important for actomyosin contraction during cytokinesis in yeast. *EMBO J.* **24**, 1440–1452 (2005).
- Stolpe, T. *et al.* In planta analysis of protein-protein interactions related to light signaling by bimolecular fluorescence complementation. *Protoplasma* **226**, 137–146 (2005).
- Marrocco, K. *et al.* Functional analysis of EID1, an F-box protein involved in phytochrome A-dependent light signal transduction. *Plant J.* **45**, 423–438 (2006).
- Atmakuri, K., Ding, Z.Y. & Christie, P.J. VirE2, a type IV secretion substrate, interacts with the VirD4 transfer protein at cell poles of *Agrobacterium tumefaciens*. *Mol. Microbiol.* **49**, 1699–1713 (2003).
- Tzfira, T., Vaidya, M. & Citovsky, V. Involvement of targeted proteolysis in plant genetic transformation by *Agrobacterium*. *Nature* **431**, 87–92 (2004).
- Loyer, A. *et al.* The plant VirE2 interacting protein 1. A molecular link between the *Agrobacterium* T-complex and the host cell chromatin? *Plant Physiol.* **138**, 1318–1321 (2005).

39. Li, J.X., Krichevsky, A., Vaidya, M., Tzfira, T. & Citovsky, V. Uncoupling of the functions of the *Arabidopsis* VIN protein in transient and stable plant genetic transformation by *Agrobacterium*. *Proc. Natl. Acad. Sci. USA* **102**, 5733–5738 (2005).
40. Lacroix, B., Vaidya, M., Tzfira, T. & Citovsky, V. The VirE3 protein of *Agrobacterium* mimics a host cell function required for plant genetic transformation. *EMBO J.* **24**, 428–437 (2005).
41. Hynes, T.R., Mervine, S.M., Yost, E.A., Sabo, J.L. & Berlot, C.H. Live cell imaging of G(s) and the beta(2)-adrenergic receptor demonstrates that both alpha(s) and beta(1)gamma(7) internalize upon stimulation and exhibit similar trafficking patterns that differ from that of the beta(2)-adrenergic receptor. *J. Biol. Chem.* **279**, 44101–44112 (2004).
42. Hynes, T.R. *et al.* Visualization of g protein beta gamma dimers using bimolecular fluorescence complementation demonstrates roles for both beta and gamma in subcellular targeting. *J. Biol. Chem.* **279**, 30279–30286 (2004).
43. Takahashi, Y. *et al.* Loss of Bif-1 suppresses Bax/Bak conformational change and mitochondrial apoptosis. *Mol. Cell. Biol.* **25**, 9369–9382 (2005).
44. Blumenstein, A. *et al.* The *Aspergillus nidulans* phytochrome FphA represses sexual development in red light. *Curr. Biol.* **15**, 1833–1838 (2005).
45. Tsuchisaka, A. & Theologis, A. Heterodimeric interactions among the 1-aminocyclopropane-1-carboxylate synthase polypeptides encoded by the *Arabidopsis* gene family. *Proc. Natl. Acad. Sci. USA* **101**, 2275–2280 (2004).
46. Ozalp, C., Szczesna-Skorupa, E. & Kemper, B. Bimolecular fluorescence complementation analysis of cytochrome P450C2, 2E1, and NADPH-cytochrome P450 reductase molecular interactions in living cells. *Drug Metabolism and Disposition* **33**, 1382–1390 (2005).
47. Szczesna-Skorupa, E. & Kemper, B. BAP31 is involved in the retention of cytochrome P450C2 in the endoplasmic reticulum. *J. Biol. Chem.* **281**, 4142–4148 (2006).
48. deVirgilio, M., Kiessens, W.B. & Shattil, S.J. Proximal, selective, and dynamic interactions between integrin alpha II beta 3 and protein tyrosine kinases in living cells. *J. Cell Biol.* **165**, 305–311 (2004).
49. Niu, T.K., Pfeifer, A.C., Lippincott-Schwartz, J. & Jackson, C.L. Dynamics of GBF1, a brefeldin A-sensitive Arf1 exchange factor at the Golgi. *Mol. Biol. Cell* **16**, 1213–1222 (2005).
50. Nyfeler, B., Michnick, S.W. & Hauri, H.P. Capturing protein interactions in the secretory pathway of living cells. *Proc. Natl. Acad. Sci. USA* **102**, 6350–6355 (2005).
51. Giese, B. *et al.* Dimerization of the cytokine receptors gp130 and LIFR analysed in single cells. *J. Cell Sci.* **118**, 5129–5140 (2005).
52. Rackham, O. & Brown, C.M. Visualization of RNA-protein interactions in living cells: FMRP and IMP1 interact on mRNAs. *EMBO J.* **23**, 3346–3355 (2004).
53. Ye, H.H., Choi, H.J., Poe, J. & Smithgall, T.E. Oligomerization is required for HIV-1 nef-induced activation of the Src family protein-tyrosine kinase, Hck. *Biochemistry* **43**, 15775–15784 (2004).
54. Schmidt, U., Richter, K., Berger, A.B. & Lichter, P. *In vivo* BiFC analysis of Y14 and NXF1 mRNA export complexes: preferential localization within and around SC35 domains. *J. Cell Biol.* **172**, 373–381 (2006).
55. Bracha-Drori, K. *et al.* Detection of protein-protein interactions in plants using bimolecular fluorescence complementation. *Plant J.* **40**, 419–427 (2004).
56. Walter, M. *et al.* Visualization of protein interactions in living plant cells using bimolecular fluorescence complementation. *Plant J.* **40**, 428–438 (2004).
57. Hackbusch, J., Richter, K., Muller, J., Salamini, F. & Uhrig, J.F. A central role of *Arabidopsis thaliana* ovate family proteins in networking and subcellular localization of 3-aa loop extension homeodomain proteins. *Proc. Natl. Acad. Sci. USA* **102**, 4908–4912 (2005).
58. Diaz, I., Martinez, M., Isabel-LaMoneda, I., Rubio-Somoza, I. & Carbonero, P. The DOF protein, SAD, interacts with GAMYB in plant nuclei and activates transcription of endosperm-specific genes during barley seed development. *Plant J.* **42**, 652–662 (2005).
59. Shimizu, H. *et al.* LIP19, a basic region leucine zipper protein, is a fos-like molecular switch in the cold signaling of rice plants. *Plant Cell Physiol.* **46**, 1623–1634 (2005).
60. Abe, M. *et al.* FD, a bZIP protein mediating signals from the floral pathway integrator FT at the shoot apex. *Science* **309**, 1052–1056 (2005).
61. Maple, J., Aldridge, C. & Moller, S.G. Plastid division is mediated by combinatorial assembly of plastid division proteins. *Plant J.* **43**, 811–823 (2005).
62. MacDonald, M.L. *et al.* Identifying off-target effects and hidden phenotypes of drugs in human cells. *Nat. Chem. Biol.* **2**, 329–337 (2006).
63. Ding, Y.H., Liu, N.Y., Tang, Z.S., Liu, J. & Yang, W.C. *Arabidopsis* GLUTAMINE-RICH PROTEIN23 is essential for early embryogenesis and encodes a novel nuclear PPR motif protein that interacts with RNA polymerase II subunit III. *Plant Cell* **18**, 815–830 (2006).
64. Wang, K.Z.Q. *et al.* TRAF6 activation of PI 3-kinase-dependent cytoskeletal changes is cooperative with Ras and is mediated by an interaction with cytoplasmic Src. *J. Cell Sci.* **119**, 1579–1591 (2006).
65. Xu, X.M. & Moller, S.G. AtSufE is an essential activator of plastidic and mitochondrial desulfurases in *Arabidopsis*. *EMBO J.* **25**, 900–909 (2006).
66. Chen, C.D., Oh, S.Y., Hinman, J.D. & Abraham, C.R. Visualization of APP dimerization and APP-Notch2 heterodimerization in living cells using bimolecular fluorescence complementation. *J. Neurochem.* **97**, 30–43 (2006).
67. Hu, C.-D. & Kerppola, T.K. in *Protein-Protein Interactions* (eds. Adams, P. & Golemis, E.) 673–693 (Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, 2005).
68. Liu, H. *et al.* Mutual regulation of c-Jun and ATF2 by transcriptional activation and subcellular localization. *EMBO J.* **25**, 1058–1069 (2006).
69. Pazhouhandeh, M. *et al.* F-box-like domain in the poliovirus protein P0 is required for silencing suppressor function. *Proc. Natl. Acad. Sci. USA* **103**, 1994–1999 (2006).
70. Cascales, E., Atmakuri, K., Liu, Z., Binns, A.N. & Christie, P.J. *Agrobacterium tumefaciens* oncogenic suppressors inhibit T-DNA and VirE2 protein substrate binding to the VirD4 coupling protein. *Mol. Microbiol.* **58**, 565–579 (2005).
71. Stains, C.I., Porter, J.R., Ooi, A.T., Segal, D.J. & Ghosh, I. DNA sequence-enabled reassembly of the green fluorescent protein. *J. Am. Chem. Soc.* **127**, 10782–10783 (2005).
72. Zhang, J., Campbell, R.E., Ting, A.Y. & Tsien, R.Y. Creating new fluorescent probes for cell biology. *Nat. Rev. Mol. Cell Biol.* **3**, 906–918 (2002).

Erratum: Design and implementation of bimolecular fluorescence complementation (BiFC) assays for the visualization of protein interactions in living cells

Tom K Kerppola

Nat. Protocols doi:10.1038/nprot.2006.201; published online 5 October; corrected online 30 November 2006.

In the version of this article initially published online, the article's page numbers should have been 1278–1286. In addition, the numbered items in Step 2 did not correspond to the format of the HTML version. In Steps 2–13, the locations of the TROUBLESHOOTING headings did not correspond to the Troubleshooting table. These errors have been corrected in the PDF version of the article.